

**UTM®**

# Universal medium for collection, transport, and preservation of viruses





Copan Universal Transport Medium (UTM®) system is intended for the collection, transport and preservation of clinical specimens containing *Viruses*, *Chlamydia*, *Mycoplasma* and *Ureaplasma*.



#### **FLOQSwabs®**



Ensure a quick, capillarity-driven sample uptake and a superior elution of the biological specimen, expanding downstream diagnostic testing capabilities.

#### **Different fill volumes and screw cap formats**



Available with a capture-cap – to dock and secure the swab shaft for easier tube handling – or an automation-friendly Pierce-able and Re-sealable PNRT™ cap\*. UTM® is available in 1 or 3mL fill volumes.

#### **Glass beads**



Three glass beads in each tube facilitate the release and dispersion of patient material and virus particles from the swab during vortexing.

#### **Skirted conical tube**



12x80mm or 16x100mm freestanding shatterproof polypropylene conical tubes ensure efficient centrifugation.

## Preservation

### UTM® Performance

Copan UTM® is **CE-IVD** and **510K cleared**, and is **compliant** with **CLSI M40-A2** standards. UTM® **preserves the viability** of all the target organisms tested **for 48 hours** at both room and refrigerated temperature.

UTM® unique formulation includes antibiotics and antimycotics to prevent overgrowth of bacterial and fungal flora without affecting viruses, chlamydia, mycoplasma, or ureaplasma viability.

According to the vast scientific literature and on the information provided by diagnostic platform manufacturers, UTM® can be used for downstream detection of:

- Nucleic acids<sup>1,2,3</sup>
- Antigens<sup>4,5</sup>
- Rickettsiae<sup>6</sup>

## FLOQSwabs®

### Cut out for everyone

FLOQSwabs® offer **variable sizes, diameters, breaking points and tip shapes to be used in plenty of applications**. This made FLOQSwabs® a well-tolerated alternative to invasive, painful, and costly collection procedures<sup>7,8</sup>

**Do you have a specific application in mind?  
Choose the right FLOQSwabs®!**



## Fields of application

### Preanalytics made different



#### *Respiratory Diseases*

Regular, minitip and flexible minitip



#### *STI & HPV*

Regular and minitip



#### *Cutaneous Infections*

Regular



#### *Veterinary*

Regular and minitip

## Laboratory

### Easy handling

Samples collected with Copan UTM-RT® can be stored at room or refrigerated temperature for 48 hours, or frozen if processing is delayed. The capture cap and the skirted tube ease the handling of the specimen, while the tubes fits most laboratory equipment. The glass beads provide a fast elution of the sample from the swab and the conical tube ensures an efficient centrifugation.



## Laboratory

### Versatile processing

UTM® can be processed both manually and with automated laboratory platforms as Copan Universe™. Copan UTM® is widely present in scientific literature and, based on information provided by the manufacturers of the diagnostic test, it can be used with:

- Antigen detection assays<sup>7,8,9</sup>
- Molecular assays<sup>10,11,12</sup>
- High-throughput systems<sup>3,13,14</sup>



#### UniVerse™

### Flexible and open solution for molecular testing sample preparation

With UniVerse™, you can automate sample preparation for molecular testing: tube decapping and recapping, vortexing, spinning, barcode identification, and liquid transfer to secondary tubes or 96-well plates. UniVerse™ handles indiscriminately different tube sizes and different shaft types without requiring to remove the swab from the tube. With its four different operational modes, UniVerse™ integrates impeccably into your lab's workflow through a 2-way LIS exchange.


Pooling samples, saving time and resources

## PodSwab™ - UTM® for pooling

PodSwab™ is a 6 ml UTM® tube paired with five nasal flocked swabs that allows for pooling up to five different individual samples in one single tube.



multiple people  
sampled



five  
FLOQSwabs®



one pooled sample



Result

NO Virus  
Detected



Virus  
Detected




### How pooling works

## The future of COVID-19 screening

If the pooled sample tests negative, all the pooled patients are considered negative – by performing one test instead of 5. Conversely, if the pooled sample is positive, each person within the pod must be individually retested.

PodSwab™ offers many operational advantages, especially in low-incidence diseases: testers can significantly increase the volume of people tested while reducing the cost per single test, in the respect of full sample and patient traceability.




UTM®





## Ordering information














Choose between different tube sizes and medium fill volumes, in bulk packs or in combination with either FLOQSwabs® or polyester fiber swabs.

### Kits of UTM® with CLASSIQSwabs™

Cat N.	Description	Pack size	Sample*
302C	<p>16x100mm tube filled with 3ml UTM® medium + 2 CLASSIQSwab™ with polyester applicator and molded breaking point</p> 	300 pieces (6 boxes of 50 kits)	Combined body site sampling

### Kits of UTM® with FLOQSwabs®

Cat N.	Description	Pack size	Sample*
305C	<p>16x100mm tube filled with 3ml UTM® medium + 1 flexible minitip FLOQSwabs® with molded breaking point</p> 	300 pieces (6 boxes of 50 kits)	naso-pharyngeal
307C	<p>16x100mm tube filled with 3ml UTM® medium + 1 minitip FLOQSwabs® with molded breaking point</p> 	300 pieces (6 boxes of 50 kits)	naso-pharyngeal, urethral, nasal, pediatric and vaginal
321C	<p>16x100 mm tube filled with 3 ml UTM® medium + 1 regular FLOQSwabs® with molded breaking point + 1 flexible minitip FLOQSwabs® with molded breaking point</p> 	300 pieces (6 boxes of 50 kits)	combined body site sampling
346C	<p>16x100mm tube filled with 3ml UTM® medium + 1 regular FLOQSwabs® with molded breaking point</p> 	300 pieces (6 boxes of 50 kits)	naso-pharyngeal, urethral, nasal, pediatric, vaginal and skin

Cat N.	Description	Pack size	Sample*
355C	<p>12x80mm tube filled with 3ml UTM® medium + 1 regular FLOQSwabs® with molded breaking point</p>  	300 pieces (6 boxes of 50 kits)	naso-pharyngeal, urethral, nasal, pediatric, vaginal, skin
358C	<p>12x80mm tube filled with 3ml UTM® medium + 1 flexible minitip FLOQSwabs® with molded breaking point</p>  	300 pieces (6 boxes of 50 kits)	naso-pharyngeal
359C	<p>12x80mm tube filled with 1ml UTM® medium + 1 regular FLOQSwabs® with molded breaking point</p>  	300 pieces (6 boxes of 50 kits)	naso-pharyngeal, urethral, nasal, pediatric, vaginal, skin
360C	<p>12x80mm tube filled with 1ml UTM® medium + 1 flexible minitip FLOQSwabs® with molded breaking point</p>  	300 pieces (6 boxes of 50 kits)	naso-pharyngeal
361C	<p>12x80mm tube filled with 1ml UTM® medium + 1 minitip FLOQSwabs® with molded breaking point</p>  	300 pieces (6 boxes of 50 kits)	naso-pharyngeal
365C	<p>12x80mm tube filled with 2ml UTM® medium + 1 flexible minitip FLOQSwabs® with molded breaking point</p>  	300 pieces (6 boxes of 50 kits)	naso-pharyngeal
366C	<p>12x80mm tube filled with 2ml UTM® medium + 1 regular FLOQSwabs® with molded breaking point</p>  	300 pieces (6 boxes of 50 kits)	naso-pharyngeal, urethral, nasal, pediatric, vaginal, skin
368C	<p>12x80 mm tube filled with 1 ml UTM® medium + 1 regular FLOQSwabs® with molded breaking point + 1 flexible minitip FLOQSwabs® with molded breaking point</p>   	300 pieces (6 boxes of 50 kits)	combined body site sampling

Cat N.	Description	Pack size	Sample*
3E076N05	12x80 mm tube filled with 6ml UTM® medium + 5 regular FLOQSwabs® with molded breaking point	300 pieces (6 boxes of 50 kits)	naso-pharyngeal



**UTM® tubes in bulk**

Cat N.	Description	Pack size	
330C	16x100mm tube filled with 3ml UTM® medium	300 pieces (6 boxes of 50 pieces)	
331C	25x90 mm tube filled with 10ml UTM® medium	90 pieces (6 boxes of 15 pieces)	
348C	16x100mm tube filled with 2,5ml UTM® medium	300 pieces (6 boxes of 50 pieces)	
349C	16x100mm tube filled with 1,5ml UTM® medium	300 pieces (6 boxes of 50 pieces)	
350C	12x80mm tube filled with 1ml UTM® medium	300 pieces (6 boxes of 50 pieces)	
353C	12x80mm tube filled with 3ml UTM® medium	300 pieces (6 boxes of 50 pieces)	

\*Suggested table. Please refer to your GLP procedures to choose the most appropriate device for the specific sampling site



## Scientific references

All the independent studies we cited in this product focus are listed here.

1. Rogers AA et al (2020) Evaluation of Transport Media and Specimen Transport Conditions for the Detection of SARS-CoV-2 by Use of Real-Time Reverse Transcription-PCR. *J Clin Microbiol* 58:e00708-20.
2. Tian J et al (2021) Identification of Viruses in Patients With Postviral Olfactory Dysfunction by Multiplex Reverse-Transcription Polymerase Chain Reaction. *Laryngoscope* 131:158-164.
3. Folgueira L et al (2019) Comparison of the Panther Fusion and Allplex assays for the detection of respiratory viruses in clinical samples. *PLoS One* 14(12): e0226403
4. Bruzzone B et al (2021) Comparative diagnostic performance of different rapid antigen detection tests for COVID-19 in the real-world hospital setting. *Int J Infect Dis* S1201-9712(21)00384-2
5. Baro B et al (2021) Performance characteristics of five antigen-detecting rapid diagnostic test (Ag-RDT) for SARS-CoV-2 asymptomatic infection: a head-to-head benchmark comparison. *J Infect* S0163-4453(21)00191-2.
6. Frickmann H et al (2013) Comparison of different media for preservation and transport of viable rickettsiae. *Eur J Microbiol Immunol* 3:194-197
7. David J. Speicher, Kathy Luinstra, Emma J. Smith, et al. Non-invasive detection of IgG antibodies from common pathogenic viruses using oral flocked swabs. *Diagnostic Microbiology and Infectious Disease*, 2020.
8. Carolynn DeByle, Lisa Bulkow, Karen Miernyk, et al. Comparison of nasopharyngeal flocked swabs and nasopharyngeal wash collection methods for respiratory virus detection in hospitalized children using real-time polymerase chain reaction. *Journal of Virological Methods*, 2012.
9. Bianco G et al (2021) Evaluation of an antigen-based test for hospital point-of-care diagnosis of SARS-CoV-2 infection. *J Clin Virol* 10438
10. Corman V et al (2021) Comparison of seven commercial SARS-CoV-2 rapid point-of-care antigen tests: a single-centre laboratory evaluation study. *Lancet Microbe*
11. Benirschke RC et al (2019) Clinical Impact of Rapid Point-of-Care PCR Influenza Testing in an Urgent Care Setting: a Single-Center Study. *J Clin Microbiol* 53:e01281-18
12. Shakya S et al (2018) Prevalence of Sexually Transmitted Infections among Married Women in Rural Nepal. *Infect Dis Obstet Gynecol*. 2018:4980396
13. Peretz A et al (2020) Mycoplasma and Ureaplasma carriage in pregnant women: the prevalence of transmission from mother to newborn. *BMC Pregnancy and Childbirth* 20:456
14. Procop GW et al (2020) A Comparison of Five SARS-CoV-2 Molecular Assays With Clinical Correlations. *Am J Clin Pathol* 155(1):69-78
15. Gomes da Costa V et al (2019) Molecular and serological surveys of canine distemper virus: A meta-analysis of cross-sectional studies. *PLoS ONE* 14(5): e0217594
16. Danilenko DM et al (2021) Antigenic and Genetic Characterization of Swine Influenza Viruses Identified in the European Region of Russia, 2014–2020. *Front. Microbiol.* 12:662028.
17. Jevsnik M et al (2020) Detection of herpes simplex and varicella-zoster virus from skin lesions: comparison of RT-PCR and isothermal amplification for rapid identification. *Diagn Microbiol Infect Dis* 97(2):115015.
18. Poljak M et al (2020) Clinical Evaluation of the cobas SARS-CoV-2 Test and a Diagnostic Platform Switch during 48 Hours in the Midst of the COVID-19 Pandemic. *J Clin Microbiol* 58: e00599-20
19. Bordi L et al (2020) Rapid and sensitive detection of SARS-CoV-2 RNA using the Simplexa™ COVID-19 direct assay. *J Clin Virol* 128: 104416.
20. Park WB, et al. Virus Isolation from the First Patient with SARS-CoV-2 in Korea. *J Korean Med Sci*, 2020 Feb 24;35(7):e84

\* The PNR cap is still under development



This document may contain product information otherwise not accessible or valid in your country. Please be aware that Copan Italia S.p.A. does take any responsibility for accessing such information which may not comply with any valid legal process, regulation, registration or usage in the country of your origin. Product clearance and availability restrictions may apply in some Countries. Please refer to Copan website ([www.copangroup.com](http://www.copangroup.com)) to view and/or download the most recent version of the brochure. This document is mainly intended for marketing purposes, always consult product insert for complete information. The use of this product in association with diagnostic kits or instrumentation should be internally validated by the user. ©2021 Copan Italia. All rights reserved. The trademarks mentioned herein are property of Copan Italia S.p.A.  
Code: JMKPF003R01



@copangroup

**Copan Italia s.p.a.**  
Via Francesco Perotti 10,  
25125 Brescia, Italy

t | f +030 2687211  
@ | [info@copangroup.com](mailto:info@copangroup.com)  
[www.copangroup.com](http://www.copangroup.com)